



Catherine B. Templeton, Director

January 30, 2015

*Promoting and protecting the health of the public and the environment*

Dear Immunization Provider:

In accordance with South Carolina Code of Laws, Section 44-29-180, and South Carolina Regulation 61-8, the 2015-2016 "Required Standards of Immunization for School Attendance" and the "Required Standards of Immunization for Day Care Attendance" are enclosed. These requirements are effective for the 2015-2016 school year.

#### **Changes for the 2015-2016 School Year**

- 11th grade has been added to the requirement for four (4) doses of diphtheria, tetanus, and pertussis.
- 1st grade has been added to the requirement for (2) doses of Varicella. A child with a positive history of the disease is considered immune and is exempt from this requirement.
- 3rd grade has been added to the requirement for three (3) doses of oral and/or inactivated Polio vaccine **with at least one (1) dose received on or after the fourth birthday.**
- 9th grade has been added to the requirement for (1) Tdap (tetanus, diphtheria, and pertussis) booster. Tdap is routinely administered at 11-12 years of age; however, a dose administered on or after the seventh birthday will meet this requirement.
- A hepatitis B third dose minimum age and interval has been added to school attendance minimum requirements (third dose received  $\geq$ 24 weeks of age and at least 16 weeks after the first dose).
- Doses documented on the immunization certificate must be valid doses according to accepted practice standards for the minimum age(s) and intervals. The "Minimum Ages and Intervals Between Doses of Routinely Recommended Vaccines" table is enclosed for your reference.

#### **General Guidance Points**

- On the day of an office visit, if a child cannot obtain all age-eligible vaccine(s) required for day care and/or school for a **medical reason**, a *medical exemption* (temporary or permanent as determined by the clinician) must be issued for the certificate to be valid. DHEC supports the provision of all Advisory Committee on Immunization Practices (ACIP) recommended vaccines according to the ACIP schedule.
- **SC State Law only allows the following exemptions to vaccine requirements:**
  - **Medical Exemption:** A valid SC Certificate of Immunization (DHEC 2740) which includes all administered vaccines to date and notes a documented medical exemption (temporary or permanent) for vaccines which are not administered for a medical reason. For temporary medical exemptions, an expiration date in the future is noted when next immunizations are due to be obtained. **THIS IS THE ONLY EXEMPTION AN IMMUNIZATION PROVIDER CAN AUTHORIZE.**
  - **Religious Exemption:** A religious exemption certificate is issued by a DHEC Health Department. The parent or guardian must sign the DHEC form in the presence of a notary.
  - **Special Exemption:** A special exemption certificate may be issued by the school to a student that has been unable to secure immunizations or documentation of immunizations already received. This exemption is only valid for 30 calendar days for the current enrollment and may be issued only once.
- A valid South Carolina Certificate of Immunization **for all enrolled children, 3 months of age and older**, must be maintained by public and private childcare facilities (as defined in Section 63-13-20 of SC Code of Laws); public, private, and parochial schools, grades kindergarten through 12<sup>th</sup> grade; and child development programs under the control of the Department of Education.
- Children enrolled in PreK or 4K programs and younger must meet Day Care Requirements, even if attendance is in a school setting.
- Children enrolled in 5-year-old kindergarten through Grade 12 must meet School Requirements.
- Children enrolled in grades 5K or greater who also attend a childcare facility (e.g., after school and/or summer program) must meet School Requirements and have a valid South Carolina Certificate of Immunization on file at both day care and school.

DHEC appreciates your continuing efforts to protect South Carolina children from vaccine-preventable diseases. Please advise your patients to receive all age-appropriate vaccines as recommended by the Centers for Disease Control and Prevention (CDC). Procedural and other questions may be directed to DHEC's Immunization Division at 803-898-0460 or 1-800-277-4687 (1-800-27-SHOTS), or [immunize@dhec.sc.gov](mailto:immunize@dhec.sc.gov).

Sincerely,

Linda J. Bell, M.D.  
State Epidemiologist

Leanne S. Bailey, RN, MPH  
Director, Division of Immunization

## 2015-2016 REQUIRED STANDARDS OF IMMUNIZATION FOR SCHOOL ATTENDANCE

Pursuant to Section 44-29-180, South Carolina Code of Laws, and South Carolina Regulation 61-8, "no superintendent of an institution of learning, no school board or principal of a school... may... enroll or retain a child or person who cannot produce satisfactory evidence of having been vaccinated or immunized so often as directed by the Department of Health and Environmental Control.... Records of vaccinations or immunizations must be maintained by the institution, school or day care facility to which the child or person has been admitted."

The Department of Health and Environmental Control has declared the following minimum immunization requirements are necessary for a child to be admitted to any public, private, or parochial school, grades five-year-old kindergarten through twelve (5K-12):

### Minimum Requirements:<sup>1,5</sup>

Vaccine for:	Grade	<i>Grade Level Requirements: Grade level requirements apply to all students entering or retained in the grades specified.</i>
<b>Diphtheria, Tetanus and Pertussis</b>	5K – 11	Four (4) doses of any combination of DTP, DT, DTaP, Td, or Tdap vaccine with at least one (1) dose received on or after the fourth birthday <sup>2</sup>
	12	Three (3) doses of any combination of DTP, DT, DTaP, Td, or Tdap vaccine with at least one (1) dose received on or after the fourth birthday
<b>Tdap Booster</b>	7-9	One (1) dose of Tdap vaccine received on or after the 7 <sup>th</sup> birthday <sup>3</sup> <b>If necessary, this dose of Tdap may be included as one of the doses needed to meet the requirement for Diphtheria, Tetanus, and Pertussis noted above.</b>
<b>Polio<sup>4</sup></b>	5K-3	Three (3) doses of oral and/or inactivated Polio vaccine with at least one (1) dose received on or after the fourth birthday
	4-12	Three (3) doses of oral and/or inactivated Polio vaccine (IPV) with at least one (1) dose received on or after the 4 <sup>th</sup> birthday <u>OR</u> four (4) doses of oral and/or inactivated Polio vaccine (IPV) before 4 <sup>th</sup> birthday (if all doses separated by at least 4 weeks) - <b>Follow CDC recommendations for students 18 years of age and older<sup>4</sup></b>
<b>Rubeola (Measles) Rubella (German Measles) Mumps</b>	5K – 12	Two (2) doses of Rubeola (Measles) vaccine with both doses received on or after the first birthday and separated by at least 4 weeks One (1) dose of Rubella (German Measles) vaccine received on or after the first birthday One (1) dose of Mumps vaccine received on or after the first birthday
<b>Hepatitis B</b>	5K – 12	Three (3) doses of Hepatitis B vaccine with the third dose received $\geq 24$ weeks of age and at least 16 weeks after the first dose
<b>Varicella</b>	5K - 1	Two (2) doses of Varicella vaccine with both doses received on or after the first birthday and separated by at least 4 weeks or a positive history of disease.
	2-12	One (1) dose of Varicella vaccine received on or after the first birthday or positive history of disease

<sup>1</sup>When issuing a South Carolina Certificate of Immunization, doses documented on the immunization certificate must be valid according to accepted practice standards for minimum age(s) and intervals.

<sup>2</sup>For unvaccinated or incompletely vaccinated children 7 years of age and older, refer to currently published ACIP Catch-Up Schedule for number of doses necessary to complete diphtheria, tetanus, and pertussis series and meet this requirement.

<sup>3</sup>Tdap is routinely administered at 11-12 years of age; however, a dose administered on or after the seventh birthday will meet this requirement.

<sup>4</sup>Routine vaccination of persons 18 years of age and older who reside in the United States is not necessary or recommended, unless in the following high risk categories: 1) traveler to areas where poliomyelitis is endemic or epidemic; 2) laboratory worker handling specimens that may contain polioviruses; or 3) healthcare worker who has close contact with patients who might be excreting wild polioviruses.

<sup>5</sup>Pursuant to Section 44-29-180, South Carolina Code of Laws, and South Carolina Regulation 61-8, students may be exempt from these immunization requirements for the following reasons:

**Medical Exemption:** The Medical Exemption section of the SC Certificate of Immunization should only be completed when a child has a permanent or temporary medical reason for exclusion from receipt of vaccine(s). The Medical Exemption section must only be completed by a licensed Physician (MD or DO) or his/her authorized representative (e.g. Physician's Assistant - PA, or Advanced Practice Registered Nurse - APRN).

**Religious Exemption:** A religious exemption may be granted to any student whose parents, parent, guardian, or person in loco parentis signs the appropriate section of the South Carolina Certificate of Religious Exemption stating that one or more immunizations conflicts with their religious beliefs. The South Carolina Certificate of Religious Exemption may only be obtained from the Department of Health and Environmental Control

**Special Exemptions:** A South Carolina Certificate of Special Exemption, signed by the school principal or his/her authorized representative, may be issued to transfer students while awaiting arrival of medical records from their former area of residence or to other students who have been unable to secure immunizations or documentation of immunizations already received. This exemption may be issued only once and is valid for only thirty (30) calendar days from the date of enrollment. Upon expiration of this special exemption, the student must present a valid South Carolina Certificate of Immunization or a valid South Carolina Certificate of Religious Exemption.

## 2015-2016 REQUIRED STANDARDS OF IMMUNIZATION FOR DAY CARE ATTENDANCE

Pursuant to Section 44-29-180, South Carolina Code of Laws, "...no owner or operator of a public or private childcare facility as defined in Section 63-13-20 may...enroll or retain a child or person who cannot produce satisfactory evidence of having been vaccinated or immunized so often as directed by the Department of Health and Environmental Control. Records of vaccinations or immunizations must be maintained by the institution, school or day care facility to which the child or person has been admitted."

The Department of Health and Environmental Control has declared the following minimum requirements are necessary to receive the final immunization certificate for day care attendance for **4K programs and younger**. Children enrolled in day care who have not yet received all required immunizations must present a valid SC Certificate of Immunization that indicates by expiration date of the certificate when the next immunization(s) are due. The child is allowed to remain in the day care **no longer than 30 days after the noted expiration date**.

**Children enrolled in grade 5K or greater AND enrolled in a childcare facility (e.g., after school and/or summer program) must meet school immunization requirements and have a valid SC Certificate of Immunization on file at the childcare facility.**

### Minimum Requirements:<sup>1, 3</sup>

Vaccine for:	Day Care Requirements:
<b>Diphtheria, Tetanus and Pertussis</b>	Four (4) doses of any combination of DTP, DT, or DTaP vaccine
<b>Polio</b>	Three (3) doses of any combination of oral or inactivated Polio vaccine
<b>Haemophilus influenzae Type b (Hib)</b>	Current, age-appropriate <i>Haemophilus influenzae</i> Type b conjugate vaccination according to the currently published immunization schedule. For children 15-59 months of age who have not yet completed age-appropriate Hib vaccination, one (1) dose of <i>Haemophilus influenzae</i> Type b vaccine at or after 15 months of age is required. <b>Hib vaccine is <u>not</u> required for children 5 years of age and older.</b>
<b>Rubeola (Measles)</b>	One (1) dose of Rubeola (Measles) vaccine received on or after the first birthday
<b>Rubella (German Measles)</b>	One (1) dose of Rubella (German Measles) vaccine received on or after the first birthday
<b>Mumps</b>	One (1) dose of Mumps vaccine received on or after the first birthday
<b>Hepatitis B</b>	Three (3) doses of Hepatitis B vaccine with the third dose received $\geq 24$ weeks of age and at least 16 weeks after the first dose
<b>Varicella</b>	One (1) dose of Varicella vaccine received on or after the first birthday or positive history of disease
<b>Pneumococcal</b>	Current, age-appropriate Pneumococcal vaccination according to the currently published immunization schedule. <sup>2</sup> For children aged 24-59 months who have not yet completed any age-appropriate pneumococcal vaccination schedule (PCV7 or PCV13), one (1) dose of 13-valent pneumococcal conjugate vaccine on or after the 2 <sup>nd</sup> birthday is required. <b>Pneumococcal conjugate vaccine is <u>not</u> required for children 5 years of age and older.</b>

<sup>1</sup>When issuing a South Carolina Certificate of Immunization, doses documented on the immunization certificate must be valid according to accepted practice standards for minimum age(s) and intervals.

<sup>2</sup>A single supplemental dose of PCV13 is strongly recommended for all children 14-59 months of age who have completed age-appropriate pneumococcal vaccination with the 7-valent pneumococcal vaccine (PCV7), but is not required for day care attendance.

<sup>3</sup>Pursuant to Section 44-29-180, South Carolina Code of Laws, and South Carolina Regulation 61-8, children may be exempt from these immunization requirements for the following reasons:

**Medical Exemption:** The Medical Exemption section of the SC Certificate of Immunization should only be completed when a child has a permanent or temporary medical reason for exclusion from receipt of vaccine(s). The Medical Exemption section must only be completed by a licensed Physician (MD or DO) or his/her authorized representative (e.g. Physician's Assistant, or Advanced Practice Registered Nurse).

**Religious Exemption:** A religious exemption may be granted to any student whose parents, parent, guardian, or person in loco parentis signs the appropriate section of the South Carolina Certificate of Religious Exemption stating that one or more immunizations conflicts with their religious beliefs. The South Carolina Certificate of Religious Exemption may only be obtained from the Department of Health and Environmental Control.

## Minimum Ages and Intervals Between Doses of Routinely Recommended Vaccines <sup>1, 2</sup>

	Vaccine and dose number	Minimum age for this dose	Minimum interval to next dose	
<b>Vaccines Required for Day Care and/or School Attendance in South Carolina<sup>1</sup></b>	Hepatitis B (HepB) -1 <sup>3</sup> HepB -2 HepB-3 <sup>4</sup>	Birth 4 weeks 24 weeks	4 weeks 8 weeks ---	
	Inactivated poliovirus (IPV)-1 <sup>3</sup> IPV-2 IPV-3 IPV-4 <sup>10</sup>	6 weeks 10 weeks 14 weeks 4 years	4 weeks 4 weeks 6 months ---	
	Diphtheria-tetanus-acellular pertussis (DTaP)-1 <sup>3</sup> DTaP-2 DTaP-3 DTaP-4 DTaP-5	6 weeks 10 weeks 14 weeks 12 months 4 years	4 weeks 4 weeks 6 months <sup>5,6</sup> 6 months <sup>5,6,7</sup> ---	
	Tetanus-diphtheria (Td)	7 years	5 years	
	Tetanus-diphtheria-acellular pertussis (Tdap) <sup>13</sup>	7 years	---	
	<i>Haemophilus influenzae</i> type b (Hib)-1 <sup>3,8</sup> Hib-2 Hib-3 <sup>9</sup> Hib-4	6 weeks 10 weeks 14 weeks 12 months	4 weeks 4 weeks 8 weeks ---	
	Measles-mumps-rubella (MMR)-1 <sup>11</sup> MMR-2 <sup>11</sup>	12 months 13 months (routine @ 4-6 years)	4 weeks ---	
	Varicella (Var)-1 <sup>11</sup> Var-2 <sup>11</sup>	12 months 15 months (routine @ 4-6 years)	See footnote <sup>12</sup> ---	
	Pneumococcal conjugate (PCV)-1 <sup>8</sup> PCV-2 PCV-3 PCV-4	6 weeks 10 weeks 14 weeks 12 months	4 weeks 4 weeks 8 weeks ---	
	<b>Recommended Vaccines<sup>1</sup></b>	Rotavirus (RV)-1 <sup>14</sup> RV-2 RV-3 <sup>15</sup>	6 weeks 10 weeks 14 weeks	4 weeks 4 weeks ---
		Hepatitis A (HepA)-1 HepA-2	12 months 18 months	6 months <sup>5</sup> ---
		Human Papillomavirus (HPV)-1 <sup>16</sup> HPV-2 HPV-3 <sup>17</sup>	9 years 9 years (+4weeks) 9 years (+24 weeks)	4 weeks 12 weeks <sup>17</sup> ---
		Meningococcal conjugate (MCV4)-1 <sup>18</sup> MCV4-2	2 years (if high risk indication) <sup>19</sup> (routine @ 11-12 years)  11 years (+8 weeks) (routine @ 16 years)	8 weeks  ---
Influenza, inactivated <sup>20</sup>		6 months	4 weeks	
Influenza, live attenuated <sup>20</sup>		2 years	4 weeks	

- <sup>1</sup> The schedule of recommended vaccines is developed and published yearly by the Advisory Committee on Immunization Practices (ACIP) and the Centers for Disease Control and Prevention (CDC). Certain ACIP-recommended vaccines are also **required** for school and/or daycare attendance in South Carolina according to the S. C. Code of Laws Section 44-20-180 and Regulation 61-8. The "Required Standards of Immunization for School Attendance" and the "Required Standards of Immunization for Day Care Attendance" are published annually by the Department of Health and Environmental Control (DHEC).
- <sup>2</sup> Combination vaccines are available. Use of licensed combination vaccines is generally preferred to separate injections of their equivalent component vaccines. When administering combination vaccines, the minimum age for administration is the oldest age for any of the individual components; the minimum interval between doses is equal to the greatest interval of any of the individual components.
- <sup>3</sup> Combination vaccines containing a hepatitis B component (Comvax, Pediarix, and Twinrix) are available. These vaccines should not be administered to infants younger than 6 weeks because of the other components (i.e., Hib, DTaP, HepA, and IPV).
- <sup>4</sup> HepB-3 should be administered at least 8 weeks after HepB-2 and at least 16 weeks after HepB-1, and should not be administered before age 24 weeks.
- <sup>5</sup> Calendar months
- <sup>6</sup> The minimum recommended interval between DTaP-3 and DTaP-4 is 6 months. However, DTaP-4 need not be repeated if administered at least 4 months after DTaP-3.
- <sup>7</sup> The fifth dose is not necessary if the fourth dose was administered at age 4 years or older.
- <sup>8</sup> Children receiving the first dose of Hib or PCV vaccine at age 7 months or older require fewer doses to complete the series.
- <sup>9</sup> If PRP-OMP (Pedvax-Hib) was administered at ages 2 and 4 months, a dose at age 6 months is not required.
- <sup>10</sup> A fourth dose is not needed if the third dose was administered on or after the 4<sup>th</sup> birthday and at least 6 months after the previous dose.
- <sup>11</sup> Combination measles-mumps-rubella-varicella (MMRV) vaccine can be used for children aged 12 months through 12 years.
- <sup>12</sup> Recommended minimum interval between doses for persons <13 years of age is 12 weeks. However, if the second dose is administered at least 28 days following the first dose, the second dose does not need to be repeated. Recommended minimum interval for persons ≥13 years of age is 4 weeks.
- <sup>13</sup> Only one dose of Tdap is recommended. Subsequent doses should be given as Td. For one brand of Tdap (Adacel), the minimum age is 11 years. For management of a tetanus-prone wound in a person who has received a primary series of a tetanus-toxoid containing vaccine, there is no minimum interval between a previous dose of any tetanus-containing vaccine and Tdap.
- <sup>14</sup> The first dose of rotavirus must be administered between 6 weeks 0 days and 14 weeks 6 days. The vaccine series should not be started after age 15 weeks 0 days. Rotavirus should not be administered to children older than 8 months 0 days, regardless of the number of doses received before that age.
- <sup>15</sup> If two doses of Rotarix are administered as age appropriate, a third dose is not necessary.
- <sup>16</sup> Bivalent HPV vaccine (Cervarix) is approved for females 10 through 25 years of age. Quadrivalent HPV vaccine (Gardasil) is approved for males and females 9 through 26 years of age.
- <sup>17</sup> The minimum age for HPV-3 is based on the baseline minimum age for the first dose (108 months) and the minimum interval of 24 weeks between the first and third doses. Dose 3 need not be repeated if it is given at least 16 weeks after the first dose (and if the intervals between doses 1 and 2 and doses 2 and 3 are maintained at 4 weeks and 12 weeks, respectively).
- <sup>18</sup> Revaccination with meningococcal vaccine is recommended for previously vaccinated persons who remain at high risk for meningococcal disease (See CDC. Updated recommendations from the ACIP for vaccination of persons at prolonged increased risk for meningococcal disease. *MMWR* 2009; 58: [1042-3]).
- <sup>19</sup> Menactra may be given as young as 2 months for high-risk children.
- <sup>20</sup> One dose of influenza vaccine per season is recommended for most people. Children younger than 9 years of age who are receiving influenza vaccine for the first time should receive 2 doses this season. See current influenza recommendations for other factors affecting the decision to administer one vs. two doses to children young than 9 years.

Adapted from: Table 1, General Recommendations on Immunization, *MMWR*, Vol. 60/No. 2, January 28, 2011.